



QUESTION WORKBOOK

Gravitational potential

Gravitational fields · Year 13

AQA 3.7.2.3 · CIE 13.4 · OCR A 5.4.4

10 questions · 30 marks

PRINT EDITION

SECTION A · Warm-up

A1 Define the gravitational potential at a point, stating where its zero is taken. [2]

A2 Calculate the gravitational potential at the surface of a planet of mass 6.0×10^{24} kg and radius 6.4×10^6 m ($G = 6.67 \times 10^{-11}$ N m² kg⁻²). [2]

A3 Explain why gravitational potential is always negative. [2]

SECTION B · Standard

B1 Calculate the gravitational potential at a point 1.0×10^7 m from the centre of the planet above (mass 6.0×10^{24} kg). [3]

- B2** A 500 kg mass is moved from the planet's surface ($r = 6.4 \times 10^6 \text{ m}$) to a point at $r = 1.28 \times 10^7 \text{ m}$. Calculate the change in gravitational potential and hence the work done. [4]

- B3** Using energy conservation, calculate the escape velocity from the surface of the planet (mass $6.0 \times 10^{24} \text{ kg}$, radius $6.4 \times 10^6 \text{ m}$). [3]

- B4** The gravitational potential changes by $1.0 \times 10^6 \text{ J kg}^{-1}$ over a radial distance of $1.0 \times 10^5 \text{ m}$. Estimate the gravitational field strength there, using $g = -\Delta V / \Delta r$. [3]

SECTION C · Stretch

- C1** Calculate the minimum energy needed to move a 1000 kg spacecraft from the surface of the planet (potential $-6.25 \times 10^7 \text{ J kg}^{-1}$) to a great distance where the potential is effectively zero. [4]

- C2** A satellite moves from an orbit at $r = 7.0 \times 10^6 \text{ m}$ to one at $r = 1.4 \times 10^7 \text{ m}$. Calculate the change in gravitational potential between the two orbits. [4]

- C3** Explain how the gravitational field strength and the gravitational potential are related through their graphs against distance. [3]