

inkMaths

ANSWER BOOK

Differentiating trig, exponentials and logs

Differentiation · Year 12–13

9 questions · 29 marks

NAVY EDITION

SECTION A · Warm-up

- A1** $\sin 2x$ gives $2 \cos 2x$, and $\cos 3x$ gives $-3 \sin 3x$. B1 B1, one for each derivative. [2]
The inner coefficient multiplies out to the front, and cosine's derivative carries the minus sign.
- A2** e^{5x} gives $5e^{5x}$, which is k times itself. $\ln x$ gives $1/x$. B1 B1, one for each derivative. [2]
The logarithm is the one entry on the shelf whose derivative leaves its own family entirely, and that is worth memorising rather than deriving under exam pressure.

SECTION B · Standard

- B1** Work termwise. $2 \sin 3x$ gives $6 \cos 3x$, and $4 \cos 2x$ gives $-8 \sin 2x$, so $dy/dx = 6 \cos 3x - 8 \sin 2x$. M1 for a correct attempt at either term, then A1 A1. Each term's inner coefficient multiplies its own front constant, so the 3 pairs with the 2 and the 2 pairs with the 4, never the other way about. [3]
- B2** $dy/dx = 5/x + 2 \cos 2x$. At $x = \pi$, $\cos 2\pi = 1$, so the gradient is $5/\pi + 2 = 3.59$. M1 [4]
A1 for the derivative, M1 for the substitution, A1 for the value. The logarithm's derivative still has an x in it, so the gradient genuinely depends on where you measure it. Working in degrees turns $\cos 2\pi$ into $\cos 6.28^\circ$ and quietly ruins the answer.
- B3** The chord gradient is $[\sin(x+h) - \sin x]/h$. Expanding the compound angle [4]
gives $[\sin x \cos h + \cos x \sin h - \sin x]/h$, and regrouping gives $\sin x (\cos h - 1)/h + \cos x (\sin h)/h$. Letting $h \rightarrow 0$ and quoting the two given limits leaves $0 + \cos x \times 1 = \cos x$. M1 for the chord gradient, M1 for the compound-angle expansion, M1 for the regrouping, A1 for the conclusion. The proof needs radians throughout, since $(\sin h)/h \rightarrow 1$ is false in degrees.
- B4** $dy/dx = 2e^{2x} + 4e^{-x}$. The -1 in the second exponent meets the -4 in front and the [3]
term turns positive, which is the sign most often lost. At $x = 0$ both exponentials equal 1, so the gradient is $2 + 4 = 6$. M1 A1 for the derivative, A1 for the value.
- B5** $\tan kx$ differentiates to $k \sec^2 kx$, so $dy/dx = 3 \sec^2 3x$. At $x = 0$, $\sec 0 = 1$ and the [3]
gradient is 3. M1 A1 A1. Since $\sec^2$ is never less than 1, this gradient never drops below 3, so $\tan 3x$ has no stationary points anywhere.

SECTION C · Stretch

-
- C1** From $x = e^y$, differentiate with respect to y to get $dx/dy = e^y$. Inverting, $dy/dx = 1/e^y$, and since $e^y = x$ this is $1/x$ as required. M1 for dx/dy , M1 for turning the rate upside down, A1 for replacing e^y by x . The final substitution is the mark students forget, leaving the answer in terms of y when the question asked for x . [3]
-
- C2** Write $a^x = e^{x \ln a}$, using $a = e^{\ln a}$. The exponent is now x times the constant $\ln a$, so the chain rule gives $\ln a \times e^{x \ln a} = a^x \ln a$. For $y = 2^x$, $dy/dx = 2^x \ln 2$, and at $x = 3$ the gradient is $8 \ln 2 = 5.55$. M1 for rewriting in base e , M1 for differentiating, A1 for the printed result, M1 for the substitution, A1 for 5.55. Treating a^x like x^n and writing $x a^{x-1}$ is the classic wrong turn. The variable is upstairs, so the power rule has no claim on it. [5]
-